

5

10 **Nicotine salts having improved taste, process for their
 preparation and their use**

15

Andreas Burgard
Margit Dörr

20

Docket 00/067 NUT

Background of the Invention

Owing to continuous consumption of cigarettes, smokers
5 are habituated to regularly taking nicotine. Because of
the disadvantageous effects of smoking on health, many
smokers are interested in giving up smoking.

This intention however fails in many cases owing to the
10 fact that the regular consumption of nicotine has already
led to a physical dependency. For slow withdrawal,
various routes have been proposed, for example applying
nicotine to the skin in the form of impregnated plasters
or supplying it orally, for example via a chewing gum
15 (US 5,488,962). Supply via chewing gum additionally has
the advantage that the oral intake compensates for the
need for sucking cigarettes.

However, nicotine has a taste which, when consumed alone,
20 is perceived as unpleasant. It is generally described as
sharp to pepper-like and bitter. This unpleasant taste
hinders the acceptance of chewing gum with added nicotine
which should be chewed regularly for withdrawal from
smoking, particularly in the initial phase. In order to
25 improve the acceptance of nicotine-containing chewing
gums or other preparations which can be consumed orally,
there is therefore the requirement for improving or
masking the unpleasant taste of nicotine.

There have been a whole series of attempts to mask the
30 unpleasant taste of nicotine. The use of flavoring
substances and taste substances such as peppermint or
strawberry flavorings has not previously led to
satisfactory masking of the nicotine taste. Only the use
35 of the inclusion compounds, for example with
cyclodextrins (WO 97/41843) is suitable to date for

inducing an improvement in taste. However, such inclusion compounds alone do not lead to sufficient taste acceptance, rather additional taste substances must be used, for example in chewing gum.

5

Therefore, the manufacturers of nicotine chewing gums have already attempted to mask the unpleasant taste of nicotine by combination with sweeteners, generally saccharin and/or acesulfame-K, which are incorporated
10 into the chewing gum base mass. The sweeteners as such, however, cannot mask the nicotine taste, so that the nicotine taste during the chewing taste becomes increasingly more clearly perceptible and becomes more clearly pronounced with the chewing process.

15

This taste impression of the abovementioned mixtures can be reproduced by simple sensory experiments with aqueous solutions, since both nicotine and the sweeteners are water soluble. For this purpose, solutions of nicotine
20 containing various sweeteners such as acesulfame-K, saccharin sodium or aspartame are prepared in water and a taste sample is compared with a nicotine-containing aqueous solution (10 mg of nicotine in 100 ml of water). The various combinations of nicotine with each of the
25 sweeteners and the corresponding taste impressions are shown in table 1.

Table 1

Taste testing of the combination of nicotine with sweeteners

5

Substances	mg of dissolved substance in 100 ml of water				
Nicotine	10	10	10	10	10
Acesulfame-K	-	12	24	12	12
Saccharin Na	-	-	-	15	-
Aspartame	-	-	-	-	15
Taste	Immedi- ately peppe r- like, rough, persis -tent unplea - santly bitter taste	Immedi- ately peppe r- like, rough, persis -tent unplea - santly bitter taste with accom- panyin g slight sweet taste	Immedi- ately pepper- like, rough, persis- tent unplea- santly bitter taste with accom- panying stronger sweet taste	Immedi- ately pepper- like, rough, persis- tent unplea- santly bitter taste with bitter and sweet after- taste	Immedi- ately pepper- like, rough, persis- tent unplea- santly bitter taste with accom- panying slight sweet taste

10

15

The results verify that the unpleasant taste of nicotine is not substantially modified by simple addition of the abovementioned sweeteners. Therefore, there is still the requirement for masking the unpleasant taste of nicotine in preparations which are taken orally, in particular in chewing gum, during the chewing process.

20

WO 99/04822 describes 1:1 salts of a sweetener and an unpleasantly tasting pharmaceutical. Salts of an anionic sweetener and nicotine are not described.

09910915 031001

WO 00/12067 describes 1:1 salts of saccharin with synthetic non-alkaloidal medicinal organic bases. Salts of an anionic sweetener and nicotine are not described.

- 5 EP-B 0 046 506 describes Vincamine saccharinate and medicaments containing this. Salts of an anionic sweetener and nicotine are not described.

10 It has now been found that nicotine, owing to its basic character, can react with anion-forming substances, which are present in the form of their corresponding acid, to give salt-like compounds.

15 Not only are salts prepared here from one molecule of nicotine and one molecule of sweetener, but also defined compounds of one molecule of nicotine and two molecules of sweetener may be prepared, in which even two different sweeteners can be used. Surprisingly, all of these compounds are distinguished by a pleasant sweet taste,
20 from which the unpleasant components of nicotine are substantially absent. They consist only of components which are used in any case, for example in nicotine-containing chewing gums, and make further working steps, such as the preparation of inclusion compounds mentioned
25 in WO 97/41843, superfluous. Thus this invention is an important advance in the production, composition and use of nicotine-containing preparations for withdrawal from smoking.

30 **Brief Description of the Invention**

The present invention thus relates to compounds or salts of nicotine) and anion-forming sweeteners (or their physiologically acceptable salts) or sweeteners which have acid character in free form, (or their physiologically acceptable salts). These compounds can also be present as

35

what are termed acid-addition salts, preferably with HCl, or other physiologically acceptable acids, for example acetic acid or sulfuric acid.

5 **Detailed Description of the Invention**

Suitable compounds for the preparation of these compounds are in principle all sweeteners or their physiologically acceptable salts, such as in particular the potassium
10 salt of acesulfame (acesulfame-K), which are able to form anions, including in particular

acesulfame and other oxathiazinone sweeteners
alitame
15 aspartame and aspartame-like di- and tripeptides
cyclamate and other sulfamate sweeteners
glycyrrhizin
neotame
saccharin
20 gluconic acid

Acesulfame (acesulfame-H) and acesulfame-K are particularly preferred.

25 A multiplicity of combinations are possible here, in particular in the case of the compounds of one molecule of nicotine and two sweetener molecules. By suitable choice of the sweeteners, in particular in compounds having two different sweeteners, surprisingly the sweet
30 taste may be modified, in particular the perception of sweetness with time, firstly with respect to masking the nicotine taste, secondly for setting an optimum overall taste impression. This is a critical advantage compared with the 1:1 compounds of 1 mol of sweetener and 1 mol of
35 nicotine.

The present invention thus also relates to a process for preparing the inventive nicotine-sweetener compounds by reacting nicotine with sweeteners in the form of their free acids (for example acesulfame-H) in a suitable solvent. Suitable solvents are preferably water and/or water-miscible solvents, for example alcohols. Sweeteners suitable for this direct reaction are preferably acesulfame, cyclamate, glycyrrhizin, gluconic acid and/or saccharin. A process variant is that the sweeteners or their salts (for example acesulfame-K) are reacted with nicotine in a solvent as described above in the presence of a physiologically harmless acid, preferably hydrochloric acid, or other suitable inorganic or organic acids, for example acetic acid or sulfuric acid, and the resultant acid addition salts (cf., for example, diagram 1) of the nicotine-sweetener compounds are isolated.

The starting materials for the preparation of the inventive compounds are commercially available or can be prepared according to methods known from the literature; for example acesulfame/acesulfame-K cf. EP-A 0 155 643.

The nicotine-sweetener compounds, salts or adducts are obtained very simply from solutions, preferably from aqueous solutions, of nicotine and the corresponding acid of the respective sweetener or, for example in a hydrochloric acid solution of the respective sweetener, as shown in the examples below, in particular for acesulfame-H. The resulting reaction solutions are freed from the solvent in a suitable manner, for example under reduced pressure. The nicotine-sweetener compounds, according to ^1H -NMR are present as 1:1, preferably as 1:2 or 1:1:1 adducts. To prepare compounds without additional salt content (no acid addition salts), preferably those sweeteners are used which are acids in the free form.

5

10

Example 1

15

20

Example 2

25


30

Example 3

35

(cf. diagram 2)

- 20 mmol (3.244 g) of nicotine are dissolved in 20 ml of water and 40 mmol (6.526 g) of acesulfame-H are added.
- 5 The reaction mixture is then concentrated under reduced pressure. A yellow amorphous solid results with 100% yield which according to $^1\text{H-NMR}$ is present as 1:2 adduct.






Diagram 2

- 5 The various nicotine-sweetener compounds in a molar ratio of 1:1 and 1:2 or 1:1:1 were subjected to sensory testing in a taste test in comparison with an aqueous nicotine solution (10 mg of nicotine in 100 ml of water). Example 4 shows the results of such testing.

10

Example 4

Nicotine-sweetener compounds in a ratio of 1:1 and 1:2 or 1:1:1

	Compound				
5	mg of dissolved adduct in 100 ml of water	1:1 nicotine-acesulfame salt 20	1:2 nicotine-acesulfame salt 31	1:1:1 nicotine-acesulfame-saccharin salt 35	1:1:1 nicotine-acesulfame - aspartame salt 66
10	Taste	Immediately sweet, after about 10 seconds, paprika-like taste, then pepper-like mild	Immediately initiating marked sweetness, persistent, initially sour, no nicotine taste, after 5 minutes: no nicotine taste	Immediately initiating weak sweetness, metallic, somewhat bitter off taste and aftertaste, after 5 minutes: no nicotine taste	Immediately initiating strong sweetness, persistent, after 5 minutes: no nicotine taste

The results verify that, in contrast to the simple mixture of free nicotine with sweeteners, the unpleasant nicotine taste can be masked by salt formation or adduct formation of nicotine with sweeteners.

Whereas in the case of the 1:1 adducts the nicotine taste can be recognized again after a relatively long residence time in the mouth, the nicotine taste in the case of 1:2 or 1:1:1 adducts of nicotine with the respective anions of the sweeteners can surprisingly even be completely eliminated, so that even after some minutes of residence time in the mouth no nicotine taste can be detected.

The inventive compounds are stable and do not decompose even during incorporation into chewing gum or other preparations suitable for withdrawal from smoking. In particular, they can be used in intermediate products or

precursor products without nicotine and sweeteners
separating. Therefore, they may be incorporated without
problem into the preparations with which they are to be
consumed, for example chewing gum, chewing tablets,
5 compressed compositions or other preparations for oral use.

The use is described by the following examples.

Example 5

10 Incorporation into chewing gum

For incorporation of the inventive compounds into chewing
gum, sugar-free chewing gum strips are produced from the
generally known ingredients chewing gum base, sugar-free
15 sweeteners, such as sugar alcohols and synthetic
sweeteners, glycerol and flavoring substances and the
inventive compounds according to generally known
technology, that is to say the ingredients are charged one
after the other into the warmed chewing gum base and are
20 incorporated uniformly. Shaping and portioning are then
carried out as usual. The ingredients correspond to those
customarily used, that is to say in the case of sugar
alcohols, for example, sorbitol, xylitol, mannitol,
maltitol, isomalt, lactitol, erythritol, mixtures of sugar
25 alcohols and sugar alcohols in syrup form, for example
sorbitol syrup and maltitol syrup. If wanted, for
additional sweetening of the chewing gums, all known
sweeteners can be used, for example acesulfame-K,
aspartame, cyclamate, saccharin, thaumatin neohesperidins
30 DC, sucralose, brazzein, neotame.

The dosage of the nicotine salts is dependent on the amount
of nicotine which is to be present in a strip of chewing
gum. In order that, for example, 2 mg of nicotine is
35 present in a chewing strip 3 g in mass, 6 mg of the
described 1:2 salt of nicotine and acesulfamic acid must be

used or else 7.61 mg of the described nicotine-acesulfamic acid-aspartame salt. Similarly, the dosage of nicotine can be increased by increasing the dosage of the nicotine salts.

5

The nicotine salts can also be incorporated into sugar-containing chewing gum formulas. For this, instead of, or together with, the sugar substitutes, the sugars customarily used, for example sucrose and glucose syrup are used. The resulting chewing gums have a pleasant sweet taste which, even in the case of relatively long chewing, is not replaced by bitter taste components.

10

Example 6

15 Incorporation into compressed compositions, tablets or chewing tablets

20

25

30

35

For incorporation of the inventive compounds into compressed compositions, sugar-free or sugar-containing compressed compositions, tablets or chewing tablets are produced with the use of sugar-free sweeteners such as sugar alcohols and synthetic sweeteners, aids, binders and flavoring substances and the inventive compounds according to generally known technology, that is to say the ingredients are homogeneously mixed one after the other and pressed to form compressed compositions or shaped to form tablets. The ingredients correspond to those customarily used, that is to say in the case of sugar alcohols, for example, sorbitol, xylitol, mannitol, maltitol, isomalt, lactitol and erythritol and mixtures of sugar alcohols. In the case of sugar-containing compressed compositions and tablets, sugars, for example sucrose, are used either exclusively or in combination with sugar alcohols. For additional sweetening of the compressed compositions and tablets, all known intensive sweeteners can be used, for example acesulfame-K, aspartame, cyclamate, saccharin,

5

The resulting compressed composition or tablet has a pleasant sweet taste which is not replaced by bitter taste components even in the event of relatively long sucking or chewing.

[illegible]